

The Riff-Shuffle Distribution is Unimodal

Stefan Gerhold*

Christian Doppler Laboratory for Portfolio Risk Management
Vienna University of Technology, Austria
sgerhold at fam.tuwien.ac.at

February 2, 2008

Abstract

We show that the probability mass function of the riff-shuffle distribution, also known as the minimum negative binomial distribution, is unimodal, but in general not log-concave.

Keywords: Discrete distribution, unimodal

2000 Mathematics Subject Classifications: Primary: 60C05; Secondary: 11K31.

The riff-shuffle distribution is defined for $0 < p < 1$ and integers $m \geq 1$ by the probability mass function

$$f_k = \binom{m+k-1}{k} (p^m q^k + q^m p^k), \quad k = 0, \dots, m-1,$$

where $p + q = 1$. If cards are taken with probabilities p and q from two decks of m cards, then the number of cards that have been taken from the remaining deck when one deck has been depleted follows this distribution [2]. Equivalently, it is the distribution of the random variable $X - m$, where X denotes the number of Bernoulli trials with individual success probability p until either m successes or m failures have been observed. This explains why it is also called minimum negative binomial distribution [3]. Johnson, Kotz, and Kemp [1] report that Uppuluri and Blot's [2] plots exhibit unimodality of the distribution. In this short note we confirm this observation.

Recall that a sequence $(g_k)_{k=0}^n$ is called log-concave if $g_k^2 \geq g_{k-1}g_{k+1}$ holds for $1 \leq k < n$. It is well known that a positive log-concave sequence is unimodal. The riff-shuffle distribution, however, provides examples of unimodal sequences that are not log-concave.

*This work was financially supported by the Christian Doppler Research Association (CDG). The author gratefully acknowledges the fruitful collaboration and support by the Bank Austria Creditanstalt (BA-CA) and the Austrian Federal Financing Agency (BFA) through CDG.

Proposition 1. *The probability mass function of the riff-shuffle distribution is log-concave for $p = \frac{1}{2}$. For $p \neq \frac{1}{2}$ and m sufficiently large, it is not log-concave.*

Proof. After cancelling the binomial coefficients, using the relation

$$\binom{u}{v} = \frac{u}{v} \binom{u-1}{v-1},$$

the log-concavity condition $f_k^2 \geq f_{k-1}f_{k+1}$ becomes

$$(p^m q^k + q^m p^k)^2 \geq \frac{k(m+k)}{(k+1)(m+k-1)} (p^m q^{k-1} + q^m p^{k-1})(p^m q^{k+1} + q^m p^{k+1}).$$

For $p = \frac{1}{2}$, this reduces to

$$1 \geq \frac{k(m+k)}{(k+1)(m+k-1)},$$

which is easily seen to be true for $k = 1, \dots, m-1$. To show the second assertion, we set $k = m-1$. After some simplifications, we see that $f_{m-1}^2 \geq f_{m-2}f_m$ is equivalent to

$$\begin{aligned} 2(2m-1)(m-1)(pq)^{2m-1} + (m-1)(p^{2m}q^{2m-2} + 2(pq)^{2m-1} + p^{2m-2}q^{2m}) \\ \geq (2m-1)(m-1) \left(\frac{p}{q} + \frac{q}{p} \right) (pq)^{2m-1}. \end{aligned}$$

The left-hand side of this inequality is $\sim 4m^2(pq)^{2m-1}$ as $m \rightarrow \infty$, whereas the right-hand side is $\sim 2(p/q + q/p)m^2(pq)^{2m-1}$. Since $p/q + q/p$ is a convex function of $0 < p < 1$ with minimum 2 at $p = \frac{1}{2}$, the right-hand side is larger than the left-hand side for $p \neq \frac{1}{2}$ and sufficiently large m . $\square$

Since the sufficient criterion of log-concavity fails us for $p \neq \frac{1}{2}$, we now prove directly that the distribution is unimodal.

Proposition 2. *The probability mass function of the riff-shuffle distribution is unimodal, with the maximum $\max f_k$ occurring for at most two (adjacent) values of k .*

Proof. For $p = \frac{1}{2}$ it is straightforward to show that $f_0 < f_1 < \dots < f_{m-2} = f_{m-1}$. Now we assume w.l.o.g. that $0 < p < \frac{1}{2}$. Defining the auxiliary function $h(x) := p^m q^x + q^m p^x$, we have

$$\frac{f_{k+1}}{f_k} = \frac{k+m}{k+1} \cdot \frac{h(k+1)}{h(k)}, \quad k = 0, \dots, m-2.$$

Therefore, $f_{k+1} \leq f_k$ is equivalent to $g(k) \geq 0$, where

$$g(x) := \frac{h(x)}{h(x+1)} - \frac{x+m}{x+1}.$$

We show that g is concave on $[0, m-2]$. The second derivative

$$\frac{d^2}{dx^2} \frac{h(x)}{h(x+1)} = \frac{(q-p)(\log p - \log q)^2 p^{m+x} q^{m+x} (p^m q^{x+1} - p^{x+1} q^m)}{(p^m q^{x+1} + p^{x+1} q^m)^3}$$

is non-positive for $0 \leq x \leq m-2$, since $(p/q)^{m-x-1} \leq 1$ implies $p^m q^{x+1} - p^{x+1} q^m \leq 0$. Thus, $h(x)/h(x+1)$ is concave. Since $(x+m)/(x+1)$ is a convex function of $x \in [0, m-2]$, the function g is indeed concave. At $x = m-2$, the function value

$$\begin{aligned} g(m-2) &= \frac{h(m-2)}{h(m-1)} - \frac{2m-2}{m-1} = \frac{p^m q^{m-2} + q^m p^{m-2}}{p^m q^{m-1} + q^m p^{m-1}} - 2 \\ &= \frac{1/p^2 + 1/q^2}{1/p + 1/q} - 2 > 0 \end{aligned}$$

is positive. From this and concavity we conclude that, if $g(l) \geq 0$ for some l , then the values $g(k)$, $l < k \leq m-2$, are positive. In other words, as soon as there is a descent $f_{l+1} \leq f_l$, the sequence $(f_k)_{k=l+1}^{m-1}$ strictly decreases. $\square$

References

- [1] N. L. JOHNSON, A. W. KEMP, AND S. KOTZ, *Univariate discrete distributions*, Wiley Series in Probability and Statistics, Wiley-Interscience [John Wiley & Sons], Hoboken, NJ, third ed., 2005.
- [2] V. R. R. UPPULURI AND W. J. BLOT, *A probability distribution arising in a riff-shuffle*, in Random counts in scientific work, vol 1: Random counts in models and structures, Pennsylvania State University Press, University Park, 1970, pp. 23–46.
- [3] Z. ZHANG, B. A. BURTNES, AND D. ZELTERMAN, *The maximum negative binomial distribution*, J. Statist. Plann. Inference, 87 (2000), pp. 1–19.